William J. Fecht, Jr., M.D.

Diplomate of the American Board of Internal Medicine / Gastroenterology

13421 Old Meridian St. Suite 201 Carmel, IN 46032

PH (317) 872-1161 TF (877) 282-5687 FX (317) 661-4657

www.indianagastro.com



NULYTELY® / GOLYTELY ® SPLIT BOWEL PREPARATION

Your efforts at cleansing your colon are essential for an accurate and safe procedure. The instructions provided below are designed to maximize the ease, safety and success of your preparation. If you believe your prep has not been effective, please inform the nurse and physician prior to the procedure.

Purchase the following at the pharmacy

- NuLytely® or GoLytely® kit (prescription provided). **Do not follow the dosing** instructions with the kit but rather those instructions below.
- Three (3) Dulcolax (bisacodyl 5 mg) tablets / pills. Available over the counter.
- Optional: Plain or aloe Baby wipes (minimizes bottom irritation)
- Optional: Desitin or A&D ointment (minimizes bottom irritation)

DAY ONE OF PREP (day BEFORE procedure)

- Eat only a clear liquid diet throughout the day. No red or purple liquids. Avoid all solid food, dairy products and juices with pulp. Examples of clear liquids can be found on the reverse side (page 2). **Drink as much fluid throughout the day.**
- Morning Mix prep according to the instructions and then refrigerate.
- 1:00 p.m. Take three Dulcolax tablets with water.
- 2:00 p.m. Begin drinking prep. Drink 8 oz every 15 30 minutes **UNTIL** you have finished <u>half</u> of the preparation. *Make sure you shake the bottle after each glass to ensure you are getting the appropriate amount of medication with each dose.*
- If moderate to severe abdominal discomfort or distension occurs, stop drinking temporarily or drink each portion at longer time intervals until symptoms disappear.
- Continue clear liquid diet.

DAY TWO OF PREP (day OF procedure)

- Drink the remaining half of your preparation. You <u>must</u> complete the preparation at least 3 hours before your procedure, so please plan accordingly.
- After completing prep, please take your necessary medications with a small amount of water.
- You may continue a clear liquid diet until 3 hours prior to your procedure.
- DO NOT TAKE ANYTHING BY MOUTH 3 HOURS PRIOR TO YOUR PROCEDURE TIME.

Please drink plenty of clear liquids throughout the day to prevent dehydration. You may have clear caffeinated beverages; however please do not make these a majority of your liquids.

CLEAR LIQUID DIET

Apple juice

Cranberry juice (okay because it is naturally red)

White grape juice

Propel fitness water (any flavor)

Gatorade (no red or purple)

Tea

Coffee. No cream or dairy products added.

Chicken or beef broth (strained)

Jell-o (no red or purple)

Popsicles (no red or purple)

You may have clear liquids with caffeine. However, please do not make it the majority of your liquids.